



Department of
Environmental
Conservation

Agriculture, Forestry, and Other Land Use

2023 NYS GREENHOUSE GAS EMISSIONS REPORT

SECTORAL REPORT #3

Kathy Hochul, Governor | Basil Seggos, Commissioner



Table of Contents

Format of This Report	1
Figures and Tables	1
Agriculture, Forestry, and Other Land Use	2
Agricultural Emissions.....	4
Livestock.....	4
Animal Feeding	4
Manure Management.....	5
Soil Management.....	7
Liming	8
Urea Application.....	8
Soil Nitrogen Management	9
Soil Organic Carbon.....	10
Net Emission Removals	12
Harvested Wood Products.....	13
Land Use	14
Forest Lands	15
Wetlands, Water Bodies, and Flooded Land	18
Cropland and Grassland	22
Settlements	23
Planned Improvements	24
Abbreviations	26
References	28

Format of This Report

This sectoral report provides a detailed explanation of methods, data, and trends for the Agriculture, Forestry and Other Lands Use (AFOLU) sector. The accounting used in this sectoral report follows the requirements of the Climate Leadership and Community Protection Act (CLCPA) and is in alignment with the 6 NYCRR Part 496 regulation, “Statewide GHG Emission Limits.” This includes the use of a 20-Year Global Warming Potential metric provided in the Intergovernmental Panel on Climate Change (IPCC) Fifth Assessment Report (IPCC 2013). The organization of this report and specific methodologies are based on the IPCC Taskforce on National Greenhouse Gas Inventories approach (or “IPCC approach”) as applied in the U.S. national greenhouse gas emissions report (IPCC 2006 and 2019, EPA 2023a). The accompanying Summary Report provides a comparison with other accounting methods, including by economic sector or using conventional accounting formats. DEC also provides emission values for all years via the Open Data NY platform.

Figures and Tables

Table SR3.1 AFOLU Agriculture Emissions, 1990-2021 (mmt CO ₂ e GWP20)	2
Table SR3.2 AFOLU Net Emission Removals, 1990-2021 (mmt CO ₂ e GWP20)	3
Table SR3.3 Livestock Emissions, 1990-2021 (mmt CO ₂ e GWP20)	4
Table SR3.4 Animal Feeding Methane Emissions, 1990-2021 (mmt CO ₂ e GWP20).....	5
Table SR3.5 Manure Management Emissions, 1990-2021 (mmt CO ₂ e GWP20)	7
Table SR3.6 Soil Management Emissions, 1990-2021 (mmt CO ₂ e GWP20)	8
Table SR3.7 Soil Nitrogen Management N ₂ O Emissions, 1990-2021 (mmt CO ₂ e GWP20).....	10
Table SR3.8 Net CO ₂ flux from soil carbon in all lands, 1990-2021 (mmt CO ₂ e GWP20)	12
Table SR3.9 Net CO ₂ Removals, 1990-2021 (mmt CO ₂ e GWP20).....	12
Table SR3.10 Harvested Wood Products CO ₂ Removals, 1990-2021 (mmt CO ₂).....	14
Table SR3.11 Land Use Net Emissions and Emission Removals, 1990-2021 (mmt CO ₂ e GWP20)	15
Table SR3.12 Forest Net Emissions and Emission Removals, 1990-2021 (mmt CO ₂ e GWP20).....	17
Table SR3.13 Wetland Net CO ₂ Removals, 1990-2021 (mmt CO ₂).....	21
Table SR3.14 Excluded Wetland CH ₄ Emissions, 1990-2021 (mmt CO ₂ e GWP20).....	22
Table SR3.15 Flooded Land Emissions, 1990-2021 (mmt CO ₂ e)	22
Table SR3.16 Cropland, Grassland, and Settlement Net Emissions and Removals, 1990-2021 (mmt CO ₂)	23

Agriculture, Forestry, and Other Land Use

This sectoral report provides information on greenhouse gas emissions and emission removals following the organizational structure of the Intergovernmental Panel on Climate Change guidance for governmental greenhouse gas inventories (e.g., IPCC 2006). For the purposes of the Climate Leadership and Community Protection Act (CLCPA), an additional distinction is made between sources covered by the statewide gross versus net emission totals, as described in the accompanying *Summary Report*. Finally, two additional sources of emissions are provided as informational items. Emissions from Soil Organic Carbon (SOC) are included in the Forest Inventory Analysis; here we report a recalculation of those emissions based on IPCC Tier 2 methods for informational purposes only. Methane emissions from wetlands are included in the Wetlands, Waterbodies, and Flooded Lands Analysis; estimated emissions are provided but are not included net totals, per IPCC accounting conventions.

In 2021, total agricultural emissions were 22.15mmt CO₂e or 41.5% higher than in 1990 (Table SR3.1). Most agricultural emissions are from livestock (92%). In 2021, agriculture represented approximately 6% of statewide emissions, when measured using CLCPA accounting. Further information on the relative contribution of the agricultural emission sources is described in the sections below. Agricultural practices also contribute to the removal of carbon dioxide from the atmosphere. These emission removals are considered a part of Land Use in the IPCC approach and are captured as “net emission removals” (Table SR3.2).

Table SR3.1 AFOLU Agriculture Emissions, 1990-2021 (mmt CO₂e GWP20)

Emission Category	1990	2005	2017	2018	2019	2020	2021
Livestock	13.56	15.15	19.55	20.09	19.75	20.15	20.33
Soil Management	2.08	2.04	1.87	1.86	1.85	1.84	1.83
Gross Total	15.65	17.18	20.42	21.95	21.59	21.99	22.15
<i>% of statewide gross total</i>	3.9%	3.7%	5.8%	5.7%	5.7%	6.4%	6.0%
Net Total	15.65	17.18	21.42	21.95	21.59	21.99	22.15
<i>% of statewide net total</i>	4.4%	4.1%	6.6%	6.4%	6.5%	7.3%	6.8%

Note: Totals may not sum due to independent rounding.

The two forms of net emission removals in New York State today are via the long-term storage of carbon in Harvested Wood Products (HWP) and the net carbon uptake and storage by various land use categories, particularly forest land (Table SR3.2). Land use and land use change drove 95% of net emission removals in 2021, but these removals have declined 14% since 1990 primarily because of forest land conversions to other land use categories.

Table SR3.2 AFOLU Net Emission Removals, 1990-2021 (mmt CO₂e GWP20)

Emission Category	1990	2005	2017	2018	2019	2020	2021
Harvested Wood Products	-1.54	-1.86	-1.56	-1.51	-1.31	-1.33	-1.45
Land Use	-31.57	-29.77	-27.87	-27.48	-27.15	-26.81	-26.71
Net Emission Removals	-33.11	-31.63	-29.50	-29.05	-29.65	-28.12	-28.04
Excluded Wetland Emissions	12.68	12.58	12.62	12.62	12.62	12.62	12.62

“+” less than 0.01mmt; “na” not applicable; Note: Totals may not sum due to independent rounding.

This inventory reports methane (CH₄) emissions from freshwater wetlands, water bodies, flooded land that has remained flooded, and land that has been converted to flooded land. Only the CH₄ emissions from the latter two flooded land categories are included in the net ‘Land Use’ totals in Table SR3.2 and contribute to the AFOLU sector’s total emissions. This is because these emissions result from human activities and land management decisions (EPA 2023a). In contrast, CH₄ emissions from freshwater wetlands and water bodies that are not considered flooded land are reported separately as ‘Excluded Wetland Emissions’ in Table SR3.2. This is in keeping with the IPCC approach, which capture anthropogenic emissions and sequestration by ‘managed lands,’ but do not count non-anthropogenic emissions from unmanaged, naturally functioning ecosystems toward inventory totals.

In the U.S., the EPA interprets flooded lands and coastal wetlands as managed lands, and therefore includes the CH₄ emissions and CO₂ sequestration fluxes attributed to these ecosystems in the national greenhouse gas inventory (EPA 2023a). However, the contributions of freshwater wetlands are only counted in the EPA’s national inventory if the wetlands are harvested for peat (EPA 2023a). DEC continues to seek feedback on whether all freshwater wetlands in New York should be considered managed lands that contribute toward the sectoral total in a similar manner to coastal wetlands. In the interim, annual CH₄ emissions from wetlands and water bodies are provided for informational purposes only, whereas the CO₂ sequestered by freshwater wetland biomass and soil carbon is counted toward the sectoral total.

These decisions affect accounting for the CLCPA net zero emission reduction goal, but they do not affect accounting for achievement of the statewide GHG emission limits established in ECL § 75-0107 and promulgated as 6 NYCRR Part 496. Finally, it should be noted that the net impact of the CH₄ emitted relative to the CO₂ sequestered by freshwater wetlands depends on the time horizon over which these are estimated. The CLCPA requires DEC to report all non-CO₂ emissions as CO₂ equivalents by using the Global Warming Potentials (GWPs) estimated over a 20-year time horizon to make the conversions. Using a 20-year rather than a 100-year GWP weights CH₄ emissions more heavily relative to an amount of CO₂ sequestered. However, protected wetland ecosystems can function over longer timescales than 20 or even 100 years, and their ability to both sequester CO₂ while also emitting CH₄ may have a net-cooling effect on climate when considered over these longer timescales (Neubauer 2014).

Agricultural Emissions

Agriculture contributes to anthropogenic emissions of carbon dioxide (CO₂), methane (CH₄), and nitrous oxide (N₂O). Livestock and soil nutrient management practices produce these emissions (Table SR3.1). Livestock sources include animal feeding (enteric fermentation) and manure storage systems. Soil nutrient management practices include liming, urea application, and soil nitrogen (N) management. These emission estimates were developed by Peter Woodbury and Jenifer Wightman (Cornell University) using IPCC methodology (IPCC 2019). Estimates of CO₂ sequestration by soils in cropland that remains cropland, and emissions that occur when other land use categories (for example, forest land) are converted to cropland were developed by the EPA (EPA, 2023a and 2023b) and are included under Land Use (Table SR3.10) in the Net Emission Removals section of this report. The agricultural industry also contributes other emissions and emission removals that are not captured in this section. For example, agriculture is a source of Energy sector emissions related to the use of fuels (*Sectoral Report #1: Energy*).

Livestock

Livestock management practices are the largest source of emissions in the agricultural sector, and they have increased 50% since 1990 (Table SR3.3). Although animal feeding is the larger source of emissions, the greatest increase in emissions is related to manure management. This trend reflects a change in policies regarding the collection and storage of manure to improve and protect water quality.

Table SR3.3 Livestock Emissions, 1990-2021 (mmt CO₂e GWP20)

Emission Category	1990	2005	2017	2018	2019	2020	2021
Animal Feeding	11.67	11.34	13.71	13.61	13.52	13.64	13.52
Manure Management	1.90	3.81	5.84	6.48	6.23	6.50	6.81
Gross Total	13.56	15.15	19.55	20.09	19.75	20.15	20.33

“+” less than 0.01mmt; “na” not applicable; Note: Totals may not sum due to independent rounding.

Animal Feeding

Ruminant animals produce CH₄ as they digest the organic carbon in their feed. Specifically, anaerobic microbes present in the animals’ digestive systems convert some of the feed into CH₄ that is then exhaled or belched by the animals.

Methodology

This analysis includes the historic and current CH₄ contribution of NY dairy cattle and non-dairy ruminant animal herds including non-dairy cattle, sheep, swine, horses, mules and asses, goats, American bison, llamas, alpacas, and deer and elk in captivity. Separate methodologies were used to estimate emissions from dairy and non-dairy animals. For non-dairy animals, herd population data were from the U.S. Department of Agriculture National Agricultural Statistics Service (NASS) (USDA 2020a) and CH₄ emissions were estimated using IPCC emission factors (Table 10.10 and 10.11, IPCC 2019). For dairy cattle, the dry matter intake (DMI) of the New York dairy herd was estimated using the number and size of cows (to calculate the energy

required for maintaining body mass), milk production, and percent milkfat (to calculate the energy required to produce the milk). The NASS provided historical data on percent milkfat, animal number, and milk production (USDA 2020a). The DMI values were used to calculate an emission factor (kg CH₄ per head per year; equation 10.21A, IPCC 2019).

Results

From 1990 to 2021, CH₄ emissions from animal feeding (enteric fermentation) increased 16% (Table SR3.4). The largest annual contributions to emissions were made by dairy cattle. Emission trends primarily reflect herd size, except among dairy cattle, whose herd sizes have decreased but whose milk production has increased over time. Annual emissions by beef cattle, for example, follow the population trends in which herd sizes increased 1990 through 2017.

Table SR3.4 Animal Feeding Methane Emissions, 1990-2021 (mmt CO₂e GWP20)

Emission Source	1990	2005	2017	2018	2019	2020	2021
Dairy Cattle	9.40	8.94	10.54	10.55	10.54	10.66	10.73
Beef Cattle	1.90	1.99	2.82	2.71	2.63	2.63	2.45
Horses	0.27	0.29	0.23	0.23	0.23	0.23	0.23
Sheep	0.07	0.06	0.06	0.06	0.06	0.07	0.06
Swine	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Goats	0.01	0.03	0.02	0.02	0.02	0.03	0.02
Deer/Elk in Captivity	ne	0.02	0.01	0.01	0.01	0.01	0.01
Bison	ne	0.01	0.01	0.01	0.01	0.01	0.01
Alpacas/Llamas	ne	+	+	+	+	+	+
Mules/Asses	+	+	+	+	+	+	+
Gross Total	11.67	11.34	13.71	13.71	13.52	13.64	13.52

“+” less than 0.01mmt; “ne” not estimated; Note: Totals may not sum due to independent rounding.

Manure Management

Manure management systems (MMS) that are used to treat and store livestock and poultry waste also emit CH₄ and N₂O. Manure stored in conditions that promote anaerobiosis will produce CH₄ (e.g., as liquid/slurry, in lagoons, ponds, tanks, or pits). Specifically, the volatile solids or organic carbon component of the manure is converted to CH₄ by certain microorganisms. In contrast, if manure is handled as a solid (e.g., in stacks or dry lots) or deposited on pasture, range, or paddock lands, it can decompose aerobically and will produce little or no CH₄. Microorganisms also produce N₂O as they act on the N compounds present in livestock and poultry waste. The series of microbial reactions that yield N₂O are known as nitrification and denitrification. They may occur directly where the waste is stored or deposited, and indirectly when the N compounds in the waste mobilize and stimulate nitrification and denitrification in another location. Two pathways produce indirect N₂O emissions. In the first, volatile manure N (mainly NH₃ and NO_x) moves through the air and is deposited on soils and the surfaces of lakes and other water bodies where it stimulates N₂O production. In the second

pathway, water in runoff and manure leachate carries N compounds into groundwater, riparian zones, ditches, streams, rivers, and estuaries that have hydrological connections to farmland. Direct and indirect N₂O emissions that are associated with manure application to soils are accounted for in the soil N management section below.

Methodology

Animal number and body weight data were gathered from the NASS (USDA 2020a). Cattle farm MMS data were gathered on DEC-regulated concentrated animal feeding operations (CAFOs) from notices of intent, change of operation, and annual compliance reports. Note that CAFOs in New York are smaller than CAFOs in other parts of the U.S. where animals may number in the tens of thousands. State permits for livestock farms in New York are based on livestock numbers alone, regardless of discharge status. Since 1999, all CAFOs operate under the no-discharge CAFO General Permit issued under the state Environmental Conservation Law (ECL) and must follow a Comprehensive Nutrient Management Plan (CNMP). Data on smaller cattle farms were gathered from the NYS Agricultural Non-Point Source Abatement and Control Grant Program (AgNPS) as well as other State data sources. Data on non-cattle MMS were gathered from livestock specialists and from annual compliance reports for non-cattle, CAFO-regulated farms.

CH₄ emissions from dairy cattle and non-dairy animals were estimated separately. IPCC emission factors were applied to each type of animal: CH₄ emissions for non-dairy animal MMS emissions were estimated using equation 10.22; dairy cattle MMS emissions were estimated using equation 10.23 (IPCC 2019). Animal waste N content was estimated using equation 10.30 and IPCC default N₂O emissions factors were applied to the fraction of manure in each type of MMS (Table 10.21, IPCC 2019).

Results

Total N₂O emissions from manure management nearly doubled between 1990 and 2021 and CH₄ emissions more than tripled (Table SR3.3, SR3.5). In 2021, total N₂O emissions from manure management were estimated to be 0.37mmt CO₂e; while in 1990 emissions were 0.19mmt CO₂e. These values include both direct and indirect N₂O emissions from manure management. Dairy accounted for 82% and 88% of N₂O emissions from the entire 1990 and 2021 livestock herds, respectively. To avoid double counting, these tables do not include emissions of N₂O from field applied manure at any time (see the Soil Nitrogen Management section). CH₄ emissions from manure management increased from 1.70mmt CO₂e in 1990 to 6.44mmt CO₂e in 2021 as a result of increased liquid storage of dairy manure.

Table SR3.5 Manure Management Emissions, 1990-2021 (mmt CO₂e GWP20)

Gas/Source	1990	2005	2017	2018	2019	2020	2021
CH₄	1.70	3.56	5.50	6.13	5.87	6.14	6.44
Dairy Cattle	1.51	3.34	5.14	5.80	5.53	5.82	6.10
Other Cattle	0.01	0.03	0.05	0.05	0.05	0.05	0.05
Swine	0.05	0.04	0.02	0.02	0.03	0.03	0.03
Chickens	0.10	0.12	0.26	0.22	0.24	0.21	0.24
Other Poultry	+	+	0.01	0.01	0.01	0.01	0.01
Horses	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Sheep	+	+	+	+	+	+	+
Goats	+	+	+	+	+	+	+
Mules/Asses	+	+	+	+	+	+	+
N₂O	0.19	0.25	0.34	0.35	0.36	0.36	0.37
Dairy Herd	0.16	0.21	0.30	0.31	0.31	0.31	0.32
Other Cattle	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Swine	+	+	+	+	+	+	+
Poultry	0.02	0.02	0.03	0.03	0.03	0.03	0.03
Sheep	+	+	+	+	+	+	+
Goats	+	+	+	+	+	+	+
Horse(mule/ass)	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Gross Total	1.90	3.81	5.84	6.48	6.23	6.14	6.80

“+” less than 0.01mmt; “ne” not estimated; Note: Totals may not sum due to independent rounding.

Soil Management

The IPCC approach include multiple emission sources under the AFOLU subcategory, “aggregated sources and non-CO₂ emissions on land.” Since the relevant sources in New York are all associated with soil amendments, we refer to this as “soil management” in this report (Table SR3.6). Other sources of emissions that are not considered relevant in New York are related to the burning of biomass as a land management practice and rice cultivation. The IPCC approach also include Harvested Wood Products in the aggregated sources subcategory but are treated as a component of net emission removals in this report (see next section).

Soil management emissions in 2021 were 4% lower than in 1990 (Table SR3.6). Most of these emissions are associated with soil nitrogen management (91% in 2021) that encompasses fertilizer application practices that result in N₂O emissions.

Table SR3.6 Soil Management Emissions, 1990-2021 (mmt CO₂e GWP20)

Gas/Source	1990	2005	2017	2018	2019	2020	2021
CO₂	0.14	0.15	0.17	0.17	0.18	0.18	0.18
Liming	0.12	0.12	0.12	0.12	0.12	0.12	0.12
Urea Application	0.02	0.02	0.05	0.05	0.06	0.06	0.06
N₂O	1.94	1.89	1.87	1.86	1.85	1.84	1.83
Soil Nitrogen Management	1.94	1.89	1.87	1.87	1.85	1.84	1.83
Gross Total	2.08	2.04	2.04	2.03	2.02	2.02	2.01

Note: Totals may not sum due to independent rounding.

Liming

Crushed limestone and dolomite are added to soils by land managers to increase soil pH. CO₂ emissions occur when these minerals react with soil acidity. Reaction rates of limestone and dolomite applications depend on soil conditions, soil type, and climate. Emissions associated with use of these minerals in industrial processes (e.g., cement production) are accounted for in *Sectoral Report #2: Industrial Processes and Product Use*.

Methodology

The average lime use over time was calculated using data from USGS (2018) and adjusted for cropland area in New York using the EPA State Inventory Tool (EPA 2023) as well as older data from the Census of Agriculture (Bureau of the Census 1989). The CO₂ emissions for each census year were calculated using the emission factors from West & McBride (2005) for limestone and dolomite and summed to estimate total CO₂ emissions due to agricultural liming.

Results

Emissions from liming of soils have fluctuated very little over the past 25 years in New York and contribute 0.12 or 0.13mmt CO₂e for each year since 1990 (Table SR3.6). Since the area of cropland changes slowly, if at all, from one year to the next, the use of lime also does not vary much.

Urea Application

Adding urea fertilizer to soils causes emission of CO₂ when the organic carbon that is integral to urea molecules is oxidized by soil microorganisms. An industrial fertilizer production process uses pressure and heat to fix CO₂ and NH₃ into urea molecules. This fixation and emission cycle is rapid, and accurately accounting for its net impact on CO₂ emissions is difficult. However, the CO₂ emitted by urea is included here to assure complete accounting for urea across sectors.

Methodology

Calendar year data on the amount of urea fertilizer applied were extracted from the SIT tool (EPA 2023). The SIT uses information on state-level fertilizer sales provided by The Tennessee

Valley Authority (1991 through 1994) and the Association of American Plant Food Control Officials. Complete conversion of all carbon in urea to CO₂ was assumed.

Results

CO₂ emissions from field application of synthetic urea increased three-fold from 0.02mmt CO₂e in 1990 to 0.06mmt CO₂e in 2021 (Table SR3.6), following increased use of urea fertilizers.

Soil Nitrogen Management

Emission sources from soil nitrogen management practices include direct and indirect N₂O emissions from managed soils, and indirect N₂O emissions from field application and management of manure. Agricultural activities that lead to direct N₂O emissions include the application of N fertilizer, the application of managed livestock manure or other organic materials such as biosolids (i.e., treated sewage sludge), the deposition of manure on soils by domesticated animals in pastures, range, and paddocks, retention of crop residues (both N-fixing legumes and non-legume crops and forages), and drainage of organic soils. Agricultural soil management activities such as irrigation, drainage, tillage, cover cropping, and fallowing of land can also influence the rates at which microorganisms produce N₂O through nitrification and denitrification and add N to soil through N fixation.

Methodology

Nitrogen from a variety of sources can be added to soil. The IPCC provides default N₂O emission factors associated with each source of added soil N (IPCC 2019). Inventoried sources of added N included synthetic fertilizer, animal manure, biosolids, and crop residues. In addition, N₂O emissions from organic soils that have been drained for agricultural activity were estimated based on the area under this type of soil and land use, and an IPCC default emission factor per unit area (IPCC 2019).

Annual estimates of synthetic N fertilizer application were from the Association of Plant Food Control Officials and The Fertilizer Institute as reported in Brakebill and Gronberg (2017) for years up until 2012. Fertilizer application was assumed to remain at 2012 levels in subsequent years. An IPCC default emission factor of 0.016 kg N₂O per kg N was used for synthetic fertilizer (Table 11.1, IPCC 2019).

Livestock manure N that was estimated for the manure management emission category (above) was assumed to be applied to soil, whether directly or after a storage period. A fraction of this applied manure N was assumed to be lost, and was subtracted from the total manure N. An IPCC default emission factor of 0.006 kg N₂O per kg N was assumed for the fraction of manure N remaining in the soil (IPCC 2019).

Data on biosolids (treated sewage sludge) applied to soil were from DEC (NYSDEC 1999 and 2018) and Baker (2016). An emission factor of 0.006 kg N₂O per kg N (IPCC 2019) was applied to an estimate of the N added to soil calculated with the total dry mass, fraction applied to soil, and the average N content of the biosolids.

The N added to soil by crop residues was estimated using crop-specific area and yields from the USDA NASS census of agriculture for barley, dry edible beans, buckwheat, corn grain, corn silage, non-alfalfa hay, alfalfa hay, non-alfalfa haylage, alfalfa haylage, hops, oats, rye, sorghum grain, sorghum silage, soybeans, sunflower, triticale, and wheat (USDA 2019). The estimated N content of specific crop residues and the emission factor of 0.006 kg N₂O per kg N were the default values provided by the IPCC (2019).

The area of drained organic soil under agricultural production was estimated for the years 1998 and 2000 by Ropel and Smith (2001) and for 2008 by the New York State Crop Reporting Service (USDA 2008). Linear interpolation was used to extend these area estimates to other years. The IPCC’s default emission factor of 13 kg N₂O-N per hectare per year was used to estimate the annual N₂O flux (IPCC 2014).

As a new approach for estimating emissions of N₂O from Histosols, the average direct emission of N₂O was estimated as the average of field measurements of the measured N₂O flux from multiple sites in NY and a site in Ohio (Duxbury & Peverly 1978, Elder & Lal 2008). Indirect emissions were calculated as part of the estimate made for mineral soils.

Results

Annual N₂O emissions from agricultural soils are estimated to be 6% lower in 2021 compared to 1990, based on available data on synthetic fertilizer sales (16% lower in recent years). Changes also reflect a decrease in N₂O emissions from daily spread of manure as dairy manure management practices have changed over time. Total N₂O emissions from soils in 2019 were largely from cropland (87%), with grassland and non-agricultural lands accounting for 8% and 5%, respectively (data not shown). In 2021, 74% of total N₂O emissions were direct, and 26% were indirect (Table SR3.7).

Table SR3.7 Soil Nitrogen Management N₂O Emissions, 1990-2021 (mmt CO₂e GWP20)

Emission Type	1990	2005	2017	2018	2019	2020	2021
Direct	1.43	1.40	1.39	1.38	1.37	1.37	1.36
Indirect	0.52	0.49	0.49	0.48	0.48	0.48	0.47
Gross Total	1.94	1.89	1.87	1.86	1.85	1.84	1.83

Note: Totals may not sum due to independent rounding; “ne” not estimated

Soil Organic Carbon

Soil organic carbon (SOC) is a major component of soil organic matter (SOM), which is a mixture of cropland residue, leaf litter, soil microbes, and products of decomposition and metabolic activity. For croplands and grasslands, SOC can be the largest stock of carbon for the ecosystem. Carbon is added via photosynthesis and the application of materials containing carbon. Carbon is lost via respiration by plants and microbes and the rate of losses can be altered by management practices such as tillage. If the gains of carbon outweigh the losses, soil carbon stocks increase (the reverse is also true). The difference between gains and losses is carbon flux, or “net” carbon emissions.

Histosols, also known as “muck” soils, are highly organic soils formed in wetlands. When they are drained, Histosols can be used as very productive cropland. However, drained Histosols also lose carbon at a much higher rate than mineral soils, and therefore have been analyzed separately from mineral soils.

This report utilizes SOC net emissions by land use type derived from FIA data, therefore these calculations are provided for informational purposes only and are not included in net totals.

Methodology

Separate methods were used to account for SOC changes in mineral soils and Histosols (drained organic soils) from 1900-2021 as described below.

For mineral soils, the IPCC Tier 1 soil organic carbon (SOC) stock change method was used (IPCC 2019, Volume 4, Chapter 2). This method makes the following two key assumptions (IPCC 2019, Volume 4, Chapter 2, page 2.32). The first assumption is that over time, SOC stocks reach a spatially averaged, stable value specific to the soil, climate, land use and management practices. The second assumption is that SOC stock change during the transition to a new equilibrium SOC occurs in a linear fashion over a period of 20 years.

Two types of activity data were used in this analysis: land use change activity data and agricultural soil management data, which includes cover cropping, tillage, and addition of organic matter. Land area data for agriculture and all other land use categories were from the United States Coastal Change Analysis Program (C-CAP). The C-CAP land area data report agriculture use in two subcategories (Cultivated, Pasture-Hay) while the USDA National Agricultural Statistics Service (USDA-NASS) land area data report agriculture use in three subcategories (Cultivated, Hay, Pasture/Other). The total area of agriculture from C-CAP was divided into subcategories using USDA-NASS data. Agricultural land can include both cropland and grassland; hay crops can be reported as either crops or grassland by different reporting bodies. For the IPCC SOC method, hay (continuous) is included in the definition of Grassland (IPCC 2019 Figure 6.1 on page 6.8). Therefore, all hay lands (both grass and alfalfa) were assigned to grassland.

For Histosols, an IPCC Tier 3 soil organic carbon (SOC) stock change method was used. New, improved estimates were made of CO₂ emissions of that use measured gas flux data from New York State and nearby regions (Ohio) in addition to an estimate based on soil subsidence. The average CO₂ emission from Histosols in New York State under conventional tillage was estimated as an average of available data from New York and Ohio (Armentano 1979, Elder & Lal 2008, Duxbury Pers. Comm.). The fraction of cropped Histosols in New York State that use no-till was provided from a personal communication from Cornell Cooperative Extension Vegetable Specialist Ethan Grundberg.

Results

Net CO₂ emissions from soil carbon in all soils in the State were 0.80 mmt CO₂e in 1990 and 0.86 mmt CO₂e in 2021 (Table SR3.8). Emissions from Histosols (organic soils) predominated with 0.80 mmt CO₂e in 1990 and 0.91 mmt CO₂e in 2021. Net CO₂ emissions from all mineral

soils were 0.00 mmt CO₂e in 1990 and -0.09 mmt CO₂e in 2021. Negative values indicate removal from the atmosphere, thus in 2021, mineral soils in the State removed 0.09 mmt CO₂e. Fluxes from all mineral soils varied over time, with some years having net emission of CO₂ from soils to the atmosphere and some years having a net removal of CO₂ from the atmosphere. Despite a 57% reduction in intensive tillage from 1990 to 2021, nearly all variation in annual SOC equilibrium values among years was due to changes in the area of cultivated mineral soils ($r^2 = 0.90$). When the cultivated area increases, emissions to the atmosphere increase, because soil carbon stocks on cultivated land are lower than those on most other land types, indicating that land use change is the predominant driver of changes in soil carbon stocks in NY.

Table SR3.8 Net CO₂ flux from soil carbon in all lands, 1990-2021 (mmt CO₂e GWP20)
Informational Purposes Only

Emission Category	1990	2005	2017	2018	2019	2020	2021
Mineral soils	-	0.03	-0.30	-0.24	-0.17	-0.09	-0.05
Drained Histosols	0.80	0.89	0.91	0.91	0.91	0.91	0.91
Total	0.80	0.92	0.62	0.68	0.74	0.82	0.86

Note: Totals may not sum due to independent rounding.

Note: Negative values are removals from the atmosphere

Net Emission Removals

The CLCPA sets three distinct emission reduction requirements, two of which relate to gross emissions to the atmosphere (Statewide GHG Emission Limits, Part 496) and a third that also relates to removals from the atmosphere, or net zero emissions. This section provides information on the two existing sources of emission removals: harvested wood products and land use. This information can be used to estimate net emission levels in the state.

Table SR3.9 illustrates the trend in net removals from 1990 through 2021 as described in more detail below. Overall net removals have decreased by 14% since 1990, but net removals associated with harvested wood products and most land use categories have remained relatively stable. For reasons discussed below, annual carbon sequestration by NY forest land has steadily declined since 1990. The excluded net emissions are CH₄ emissions from wetlands and water bodies.

Table SR3.9 Net CO₂ Removals, 1990-2021 (mmt CO₂e GWP20)

Emission Category	1990	2005	2017	2018	2019	2020	2021
Harvested Wood Products	-1.54	-1.86	-1.56	-1.56	-1.51	-1.31	-1.33
Land Use	-32.18	-30.96	-28.46	-28.12	-27.79	-27.88	-27.54
Net Emission Removals	-33.73	-32.82	-30.03	-29.68	-29.30	-29.19	-28.87
Excluded Net Emissions	12.68	12.58	12.62	12.62	12.62	12.62	12.62

Note: Totals may not sum due to independent rounding.

Harvested Wood Products

Most timber harvested from forest land in New York is used to produce harvested wood products (HWPs), which include, for example, sawn wood used in housing construction and paper. HWPs sequester CO₂ for a duration that depends on the lifespan of the product and the rate at which it is converted back to CO₂ through decay in landfills or waste combustion.

Emission estimates for this section of the report were developed in conjunction with Robert Malmshaimer, HakSoo Ha, Timothy Volk, Tristan Brown, Danielle Kloster, Jenny Frank, Theodore Koch, Nehan Naim, and Obste Therasme (State University of New York, College of Environmental Science and Forestry) and the DEC Division of Lands and Forests Forest Utilization Program.

Methodology

The accounting approach taken includes carbon storage by sawtimber-based HWPs but does not include pulp logs used to produce paper or HWPs used to produce other products with shorter useful lives. Sourcing includes trees that are (1) grown in New York and consumed outside New York (NY/Out), (2) grown outside New York and consumed in New York (Out/NY), or (3) grown and consumed within NY (NY/NY). This approach captures out-of-state trees that are imported for use within New York, as well as trees that are grown within New York and exported for out-of-state for use.

Mill survey data collected by DEC, known as the Timber Products Output (TPO) data, were used to estimate annual board feet of sawtimber-based HWPs used between 1999 and 2021, and were then converted to carbon storage. Trends from this data were extrapolated back through time to produce an estimate of the carbon stored in wood products harvested from 1990 to 1998, a period before the data collection program had started. This method converted board feet of lumber to carbon storage assuming a density (oven-dry mass / air-dry volume) of 0.56 metric tonnes per cubic meter for hardwood and 0.49 metric tonnes per cubic meter for softwood, and carbon fractions of 0.5 for both types of wood. Using this dataset as a proxy for HWPs may overestimate carbon storage in HWPs because conversion of sawtimber to HWPs is not complete and does not account for mill residuals that may be burned to produce energy. The dataset may underestimate carbon storage by non-lumber HWPs such as paper and cardboard that are produced from pulpwood and chips. Finally, this component of the inventory does not explicitly model the decay or combustion of HWPs in landfills and other solid waste facilities, although the solid waste sector analysis in *Sectoral Report #4: Waste Report* of this inventory includes a decay model for organic carbon in landfills.

There may be other emissions associated with production of timber including from fertilizer application or the energy used to harvest, process, and transport the harvested materials. These emissions are captured in relevant agriculture and energy emission sectors. The impact of timber harvesting on carbon sequestration by forested lands is captured in the land use section below.

Results

Annual trends in CO₂ removal by HWP roughly correspond with those in new housing construction in the Northeast, peaking in 2001 at -1.97mmt CO₂, declining to a minimum of -1.17mmt CO₂ per year in 2009, recovering to an average of -1.57 from 2015 to 2018. The drop in 2019 and 2020 to -1.31 and -1.33mmt CO₂, respectively, may reflect changes in mill survey response rates during the COVID-19 pandemic.

Table SR3.10 Harvested Wood Products CO₂ Removals, 1990-2021 (mmt CO₂)

Origin/Destination	1990	2005	2017	2018	2019	2020	2021
NY/NY	-1.09	-1.21	-1.07	-1.05	-0.94	-0.92	-1.09
NY/Out	-0.42	-0.56	-0.32	-0.29	-0.26	-0.25	-0.25
Out/NY	-0.03	-0.09	-0.17	-0.17	-0.11	-0.16	-0.11
Net Emission Removals	-1.54	-1.86	-1.56	-1.51	-1.31	-1.33	-1.45

Note: Totals may not sum due to independent rounding.

Land Use

The analysis of net emission removals by land in New York included an assessment of both land use and land use change for each year between 1990 and 2021. The annual fluxes of greenhouse gases into and out of natural and managed lands depends mainly on the growth and decay of trees, plants, and other organisms that can remove CO₂ from the atmosphere and sequester it as organic carbon. Certain landscapes, such as forests, sequester and store more organic carbon per unit area than other lands.

The six categories of land use that were considered here are forest land, wetlands, cropland, grassland, settlement, and other lands, following the IPCC approach (IPCC, 2019). Quantifying the net emission removals through land use and land use change requires accounting for the differences in carbon sequestration and greenhouse gas emissions per unit area among land use and surveying changes in the areal extent of land use categories. Land parcels that have remained under the same category of land use for 20 years or more are classified as forest remaining forest, grassland remaining grassland, cropland remaining cropland, etc. Lands that have been converted from one land use category to another within 20 years or more are classified based on their land use categories before and after conversion (e.g., cropland converted to forest, forest converted to grassland, etc.).

The EPA has made available net emissions and emission removals estimates for cropland, grassland, and settlements in New York from 1990 through 2021 (EPA 2023b and Table SR3.11). These estimates are now included along with emission removals by forest land and wetlands. This aligns this inventory with the IPCC's guidance that emissions or emission removals associated with land that remains under a particular land use category (e.g., forest land) should be added to emissions or emission removals associated with any conversions to that land use category (e.g., cropland to forest land, grassland to forest land, settlement to forest land, etc.). Note that unlike the other land use categories, the EPA's cropland estimates

reflect soil carbon stock changes and not carbon storage in biomass, litter, or dead organic matter on that cropland (EPA 2023a and 2023b).

Table SR3.11 Land Use Net Emissions and Emission Removals, 1990-2021 (mmt CO₂e GWP20)

Land Type	1990	2005	2017	2018	2019	2020	2021
Forest land	-31.34	-29.45	-27.85	-27.60	-27.33	-27.07	-26.80
Forest converted to other land	0.27	0.26	0.26	0.26	0.26	0.26	0.26
Wetlands	-0.89	-0.73	-0.81	-0.81	-0.81	-0.81	-0.81
Flooded land	2.10	2.09	2.09	2.09	2.09	2.09	2.09
Cropland	-0.10	0.07	0.46	0.54	0.58	0.48	0.54
Grassland	-0.38	-0.98	-0.27	-0.27	-0.24	-0.31	-0.28
Settlement	1.67	1.69	1.75	1.75	1.75	1.74	1.75
Settlement trees	-3.52	-3.91	-4.09	-4.08	-4.08	-4.26	-4.28
Net Emission Removals	-32.18	-30.96	-28.46	-28.12	-27.79	-27.88	-27.54

Forest Lands

New York is among the most forested states in the nation with roughly 19 million acres of forest land, covering over 60% of the land area. This is in large part why lands in New York remove more CO₂e than they emit on an annual basis. Specifically, forest land that has remained forest land removes additional CO₂ as organic carbon every year, equivalent to -26.80mmt CO₂ in 2021 (Table SR3.12). The conversion of other land use categories to forest land removed an additional -2.15mmt CO₂ in 2021 because forest land removes more CO₂ per unit area than any other land use category.

Methodology

This report uses state data from the EPA and U.S. Forest Service (Forest Service) on annual forest CO₂e emissions and sequestration (Domke et al. 2023; EPA 2023b; Walters et al. 2022). Additional input on this analysis was received from Colin Beier (State University of New York, College of Environmental Science and Forestry) as part of ongoing work (see the Planned Improvements section). The annual Forest Service analysis assesses the sequestration of CO₂ as organic carbon in five reservoirs: aboveground biomass, belowground biomass, dead wood, litter, and soil organic carbon. The rate of change in the organic carbon density of these reservoirs is assessed by sampling specific field plots over time and measuring biomass, dead wood, and litter. Soil organic carbon is modeled using latitude, elevation, precipitation, temperature, and moisture index data (Domke et al. 2017) as well as remote sensing information from the National Agriculture Imagery Program (NAIP). The assessment implicitly incorporates any losses of organic carbon density and CO₂ emissions from fires, forest cutting, insects, disease, and weather events (Coulston et al. 2015). The impact of land use change on forest land is evaluated using measured annual rates of transitions among cropland, grassland, settlement, forest land, wetlands, and other land use categories. Observed conversions of field plots from one land use category to another are used to quantify the impact of the conversions on organic carbon stocks in the absence of other data. Conversions of forest to cropland or

grassland include changes in soil organic carbon stocks, aboveground and belowground biomass, dead wood, and litter carbon stocks. 'Forest land converted to other land' is currently an orphan land use conversion type that is estimated by the U.S. Forest Service and will be combined into an 'other land' use category when a complete estimate becomes available. Trees located in settlements that remain settlements are reported as "settlement trees."

The Forest Inventory and Analysis (FIA) program published a new model in 2023 for predicting tree cubic-foot volume, biomass, and carbon attributes. This National Scale Volume and Biomass Estimators system provides a more consistent and accurate accounting of structural components of trees across the U.S. for total tree cubic-foot volume, biomass, and carbon (Westfall et al. 2023). The new model has informed the FIA data used in this inventory, and thus the current inventory reflects these changes in forested lands, land conversions, land use, and settlement trees.

Achieving a wall-to-wall assessment of carbon sequestration and emissions by the different land use categories will require additional work to reconcile the area considered "forest" according to the Forest Service's inventory, and the areas that may be mapped as forest by satellite-based land-cover products (such as those used to assess wetlands in this report). The land use definition of forest land that is used here may include some harvested areas that are replanted or left to regenerate and forested areas that lose biomass through forest fires or other disturbances. Land-cover maps may classify these areas into categories other than forest (Coulston et al. 2015), causing overlaps in coverage. Furthermore, some forested areas that do not meet the Forest Service's definition of forest are not counted in this inventory, including areas that are too small (less than 120 feet wide and 1 acre in area with at least 10% cover; Oswalt et al. 2019) and agroforestry systems (EPA 2023a).

A subset of land defined as forest land in the Forest Service's inventory is also wetland. These lands were excluded from the wetlands component of this report to avoid double-counting (an average of -3.93mmt CO₂ sequestered per year from 1990 to 2021). However, forested wetlands also have the potential to produce CH₄ and N₂O emissions as well as sequester CO₂. The Forest Service inventory does not currently include these emissions, so they are also omitted in the forest land estimates in this report, but they are discussed in the wetlands section (below).

Results

The forest land use category includes emissions and emission removals associated with forest land remaining forest land and conversions of other land use categories to forest land. In the previous inventory, forest land conversions to other land use categories were also included in the forest land use category. Although these emissions are still reported in Table SR3.12, they are counted implicitly among the other land use categories as conversions. For inventory purposes, forest land conversions are now split up among the other land use categories so that, for example, emissions associated with forest land converted to cropland is now considered a part of the cropland category's emissions. Most net emission removals were by forest land that remained forest land, but an additional -2mmt CO₂ per year was also removed through the

conversion of cropland, grassland, settlements, wetlands, and other lands to forest land. However, the emission removals by conversions of the other land use categories to forest land were more than offset by over 3mmt CO₂ of emitted annually as forest land is converted to other land use categories. Finally, emissions of CH₄ and N₂O from forest fires were estimated and reported by the Forest Service and are accounted for in the net emission removals attributed to forest land (Walters et al. 2023).

Table SR3.12 Forest Net Emissions and Emission Removals, 1990-2021 (mmt CO₂e GWP20)

Land Use Change	1990	2005	2017	2018	2019	2020	2021
Forest Remaining Forest	-31.35	-29.45	-27.85	-27.60	-27.33	-27.07	-26.80
Land Converted to Forest	-2.15	-2.12	-2.09	-2.09	-2.09	-2.09	-2.09
Forest Converted to Land	3.23	3.35	3.47	3.46	3.46	3.47	3.46
Forest Fire	nd	nd	nd	nd	nd	+	nd
Net Emission Removals	-30.27	-28.22	-26.47	-26.23	-25.96	-25.69	-25.43

“nd” no data; “+” positive values less than 0.01mmt

Forest Land Remaining Forest Land

Forest land remaining forest land sequestered far more CO₂ than any other land use category (-29.45mmt CO₂ sequestered in 2005 and -26.80mmt CO₂ sequestered in 2021). The declining trend in annual CO₂ sequestration is due in part to the conversion of forest land to other land use categories (3.46mmt CO₂ emitted in 2021 and an average of 3.46mmt CO₂ emitted per year between 2005 and 2021), which was not matched by the rate at which CO₂ was sequestered as other land use categories converted to forest land (-2.09mmt CO₂ sequestered in 2021 and an average of -2.10mmt CO₂ per year sequestered between 2005 and 2021). Future inventories will include an assessment of the impact of interannual changes in the rate at which forest land that remains forest land sequesters CO₂ and stores it as organic carbon.

Land Converted to Forest Land

Cropland conversion to forest land sequestered an average of -0.61mmt CO₂ per year from 2005 to 2021. Settlement conversions to forest land sequestered an average of -1.02mmt CO₂ per year from 2005 to 2021. Wetlands converted to forest land sequestered an average of -0.28mmt CO₂ per year from 2005 to 2021. Other lands converted to forest land sequestered -0.20mmt CO₂ per year from 2005 to 2021. Grassland converted to forest sequestered an average of -0.02mmt CO₂ per year from 2005 to 2021.

Forest Land Converted to Land

Conversion of forest land to cropland resulted in an average of 1.33mmt CO₂ emitted per year between 2005 and 2021. Conversion to settlements resulted in an average of 1.37mmt CO₂ emitted per year between 2005 and 2021. Smaller emissions were caused by conversion to wetlands (an average of 0.41mmt CO₂ emitted per year between 2005 and 2021) and other lands (an average of 0.34mmt CO₂ emitted per year between 2005 and 2021). There were no emissions attributed to grassland conversions between 2005 and 2021.

Wetlands, Water Bodies, and Flooded Land

Emissions and emission removals attributed to wetlands and CH₄ emissions from other water bodies are reported here along with EPA's estimates of CO₂ and CH₄ emissions from land flooded as a direct result of human interventions such as dam and pond construction (EPA 2023a). Wetlands contribute most of the emissions and emission removals estimates in this section. Wetlands, like forest land, sequester CO₂ and store it as organic carbon in plant biomass and soils. Unlike forest land, whose area is estimated by the Forest Service according to a land use definition, the areal extent of wetlands is estimated here using land cover observations. All New York wetlands are inventoried here, even those that do not directly undergo management activities, such as peat harvesting. New York has both coastal wetlands, which are exposed to tidal cycles and are often adapted to brackish and saltwater inundation, and upland wetlands, which are not tidal and whose soils are saturated with freshwater.

Vegetated wetlands are particularly effective at sequestering CO₂ as organic carbon and storing it in their soils. This is because water-saturated soils tend to be low-oxygen soils, and oxygen scarcity slows the rate of decay that would normally convert organic carbon back into CO₂ (IPCC 2014). Wetland soils can also be important sources of CH₄ and N₂O. Freshwater wetlands are an important source of CH₄ whereas CH₄ production is suppressed in tidal wetlands that are inundated by brackish and salt water above a certain salinity threshold (Holmquist et al 2019; IPCC 2014). CH₄ emissions from freshwater wetlands are provided for informational purposes and are not included in net totals. N₂O production may occur in either type of wetland and is stimulated by the discharge of nitrogen (e.g., nitrate and ammonium) from the watershed, as well as atmospheric deposition of nitrogen compounds generated by fossil fuel combustion or volatilized when nitrogen fertilizers are applied to soil.

Methodology

The distinction between wetlands that are inundated by freshwater versus brackish to saltwater is made here using land-cover maps from satellite images that are produced by the Coastal Change Analysis Program (CCAP) within NOAA. These maps classify wetlands as either estuarine or palustrine, where palustrine wetlands are inundated by water with a nominal salinity that is less than 0.5 parts per thousand (ppt). Estuarine wetlands are assumed to produce no CH₄, whereas the emission factor for CH₄ from palustrine wetlands is assumed to be 157 kg CH₄ per hectare per year (see Section 4.3.1.2 and Annex 3A.3 in IPCC 2014). The CCAP maps further classify estuarine and palustrine wetlands as forested, scrub and shrub, or emergent vegetation, for a total of six wetlands land-cover categories. Annual changes in the areal extent of the six wetlands categories were estimated by comparing consecutive updates of the CCAP land-cover maps. Beginning in 1996, the maps have been issued approximately every 5 years, with the most recent update issued in 2016. The areas of wetlands that remain wetlands as well as conversions of other land-cover types to and from wetlands has been interpolated to cover years between map updates. After 2016, wetland areas have been held constant and will be updated with appropriate interpolations once the next CCAP land-cover classification map is made available.

Wetland carbon sequestration is assessed by the following organic carbon reservoirs: soil carbon and aboveground and belowground biomass. Annual CO₂ fluxes into soil carbon are geometric means of soil and sediment core measurements made using lead-210 and retrieved from publications and public databases by the Smithsonian's Environmental Research Center.¹ The carbon sequestration by aboveground biomass and belowground biomass are the same values used in the national greenhouse gas inventory (EPA 2023a). Aboveground wetland biomass is assumed to sequester -3.05 to -3.17mt of carbon per hectare per year. Belowground biomass sequesters -6.44 to -6.54mt carbon per hectare per year in estuarine wetlands, and -3.57 to -3.65mt carbon per hectare per year in palustrine wetlands. For conversions of other land-cover types to and from wetlands, the impact of biomass in the other land-cover types is captured here but changes in CO₂ sequestration in soil carbon or soil carbon stock changes are not captured.

Areas that are classified as palustrine forested wetlands in the CCAP land-cover maps are assumed to have been counted as forest land in the Forest Service's national inventory report data (Domke et al. 2023, Walters et al. 2023). Therefore, they have not been included in the CO₂ sequestration calculations for palustrine wetlands, but their CH₄ emissions are included here. No land areas classified as estuarine forested wetlands by CCAP are included in the Forest Service's inventory because data are needed to quantify the impacts of forestry activity on soil carbon stocks in forested coastal wetlands (Crooks et al. 2018, Lisa Schile-Beers personal communication). Therefore, these areas are included in calculations of CO₂ sequestration by estuarine wetlands. The CCAP maps also resolve 19 other land-cover classes that are grouped to approximate the other five IPCC land use categories in order to quantify conversions of wetlands to and from other types of land cover (IPCC 2019).

Seagrass is included in the calculation of CO₂ sequestration by estuarine wetlands. The areal extent of seagrass in NY waters is assumed to reflect a value reported in the 2009 Final Report of the New York State Seagrass Task Force.² In the absence of regular, state-wide survey data, this area is held constant from 1990 to 2021 and multiplied by the seagrass soil carbon sequestration fluxes reported by Salinas et al. (2020).

The USGS National Hydrography Dataset (NHD) maps (USGS 2016) were used to estimate CH₄ emissions from lakes, ponds, and other water bodies in New York. The water bodies were grouped into 7 size classes (<0.001 km² to >100 km²) and their areas multiplied by the size-specific CO₂ and CH₄ emission fluxes according to Holgerson and Raymond (2016). Emissions from water bodies that cross state boundaries (Lake Erie, Lake Ontario, Lake Champlain, and the Allegheny Reservoir) are scaled to the proportions of the water bodies' total areas that fall within the state's political boundaries. Mapped stream and river features are not included in this estimate pending further method development. There is a small amount of overlap between water bodies mapped in the NHD and the emergent wetland areas mapped by CCAP. There are also areas of open water mapped by CCAP that are not mapped by NHD. Additional work is

¹ <https://github.com/Smithsonian/Coastal-Wetland-NGGI-Data-Public>

² https://www.dec.ny.gov/docs/fish_marine_pdf/finalseagrassreport.pdf

necessary to determine the extent and impact of these coverage differences between the two data sets. Emission factors for CH₄ ranged from 3.5 to 133.5kg CH₄ per hectare per year for water bodies with areas greater than 100 km² to water bodies with areas smaller than 0.001 km², respectively (Holgerson and Raymond 2016). CO₂ emissions were not estimated to avoid double counting CO₂ derived from organic carbon sourced from other land use categories.

The EPA's state estimates of CH₄ and CO₂ emissions from land flooded as a result of direct human interventions includes reservoirs, canals and ditches, constructed freshwater ponds, and inundation areas (EPA 2023a and 2023b; IPCC 2019). The emissions calculations distinguish between land that has been flooded for more than 20 years (flooded land remaining flooded land) and land that has been flooded for less than 20 years (land converted to flooded land) because emission factors are higher for recently flooded land. Emission factors were also split between the state's two climate zones, warm temperate moist and cool temperate. Flooded land was identified and areas estimated using NHD maps, the National Lakes Assessment (EPA 2023c), the National Inventory of Dams (USACE 2021), and the Navigable Waterways dataset (DHS 2022). Note that there may be overlap in the emissions estimated for water bodies, which include reservoirs and ponds, and the subset of reservoirs and ponds that are considered flooded land by the EPA. For flooded land remaining flooded land, surface emission factors of 54 or 80kg CH₄ per hectare per year were applied to reservoirs in cool temperate or warm temperate moist zones, respectively, along with a 9% downstream emission factor (IPCC 2019). CO₂ emissions were not estimated for flooded land remaining flooded land to avoid double counting CO₂ derived from organic carbon sourced from other land use categories. For land converted to flooded land, surface emission factors of 128 or 85kg CH₄ per hectare per year were applied to reservoirs in warm temperate moist or cool temperate climate zones, respectively, along with a 9% downstream emission factor. CO₂ surface emission factors applied to freshwater ponds and reservoirs were 5.4 or 3.7mt CO₂ per hectare per year in the warm temperate moist or cool temperate climate zones, respectively. The same surface emission factor of 183kg CH₄ per hectare per year was applied to all constructed freshwater ponds, regardless of their age or climate zone. The highest CH₄ emission factor (416kg CH₄ per hectare per year) was applied to canals and ditches in both climate zones (IPCC 2019).

Results

The wetland category is broken into four types of land use and land use change, as described above (Table SR3.13). Palustrine wetlands were responsible for 94% of annual removals by wetlands in 2021. However, this accounting does not include CH₄ emissions from palustrine wetlands (Table SR3.14).

Table SR3.13 Wetland Net CO₂ Removals, 1990-2021 (mmt CO₂)

Land Use Change	1990	2005	2017	2018	2019	2020	2021
Estuarine Wetlands Remaining Estuarine Wetlands	-0.09	-0.08	-0.08	-0.08	-0.08	-0.08	-0.08
Palustrine Wetlands Remaining Palustrine Wetlands	-1.18	-1.13	-1.22	-1.22	-1.22	-1.22	-1.22
Land Converted to Wetlands	0.38	0.43	0.49	0.49	0.49	0.49	0.49
Wetlands Converted to Water	nd	0.06	nd	nd	nd	nd	nd
Net Emission Removals	-0.89	-0.73	-0.81	-0.81	-0.81	-0.81	-0.81

“nd” no data

Wetlands Remaining Wetlands

Palustrine wetlands sequester more CO₂ (-1.22mmt CO₂ in 2021) than estuarine wetlands (-0.08mmt CO₂ in 2021), mainly because palustrine wetlands occupy more land in NY. Forested palustrine wetlands sequester more CO₂ than any other type of wetlands (-3.91mmt CO₂ in 2021) due to their larger area and capacity to form biomass. However, to avoid double counting, these removals are not included in the net emission removals by wetlands in Table SR3.13 because they are included in the forest land areas contributing net emission removals in Table SR3.12. Estuarine wetlands include -0.02mmt CO₂ sequestered per year as soil carbon accumulated by seagrass meadows in NY waters and -0.06mmt CO₂ sequestered per year by estuarine wetlands as biomass and soil carbon (Table SR3.13). Estimated CH₄ emissions from palustrine wetlands and water bodies are 0.14mmt CH₄ per year (11.99mmt CO₂e) and 0.007mmt CH₄ (0.63mmt CO₂e) per year, respectively (Table SR3.13). Forested palustrine wetlands contribute 0.11mmt CH₄ (9.14mmt CO₂e) per year to the palustrine wetlands total, and this is reported here because the Forest Service does not currently include this CH₄ source in its forest inventory (Grant Domke, personal communication). The CH₄ emissions reported in this section are for informational purposes only, as discussed above.

Wetlands Converted to Water

Sea level rise, coastal subsidence and erosion can convert vegetated coastal wetlands to open water. When this happens, some of the organic carbon stored in wetland biomass and soil can be converted back to CO₂. Catastrophic coastal erosion can cause the release of many years of accumulated soil carbon stock (Crooks et al. 2018, Holmquist et al. 2018). In New York, conversion of wetlands to water resulted in the emission of 0.06mmt CO₂ per year between 2002 and 2010 but were less than 0.01mmt CO₂ per year from 1990 to 2001 and from 2011 to 2021.

Lands Converted to Wetlands

Conversion of forest land to wetlands emitted 0.49mmt CO₂ in 2021 (Walters et al. 2023). Conversion of other lands to wetlands contributed an average of 200 metric tons of CO₂ per year from 1990 to 2021.

Excluded Wetland Emissions

As indicated in the introduction to this report, this assessment omits CH₄ emissions by freshwater wetlands and water bodies (Table SR3.14) from total net removals by wetlands. If these emissions were included, they would be subtracted from net emission removals attributed to the land use categories above (Table SR3.13). These estimates are also preliminary, and these emissions are not included in the U.S. national greenhouse gas inventory (EPA 2023a).

Table SR3.14 Excluded Wetland CH₄ Emissions, 1990-2021 (mmt CO₂e GWP20)
Informational Purposes Only

Emission Source	1990	2005	2017	2018	2019	2020	2021
Palustrine Wetlands	12.05	11.95	11.99	11.99	11.99	11.99	11.99
Water Bodies	0.63	0.63	0.63	0.63	0.63	0.63	0.63
Excluded Net Emissions	12.68	12.58	12.62	12.62	12.62	12.62	12.62

Flooded Land

Emissions of CO₂ and CH₄ from land flooded through direct human intervention are included in total net removals (Table SR3.15). These estimates are included in the U.S. national greenhouse gas inventory (EPA 2023a) and the net emissions of this inventory. Nearly all emissions were CH₄ (2.09mmt CH₄ in 2021) and reservoirs were responsible for most of the emissions (54% from 1990 to 2021).

Table SR3.15 Flooded Land Emissions, 1990-2021 (mmt CO₂e)

Type	1990	2005	2017	2018	2019	2020	2021
CH₄	2.09	2.09	2.09	2.09	2.09	2.09	2.09
Reservoirs	1.14	1.14	1.14	1.14	1.14	1.14	1.14
Constructed Ponds	0.92	0.92	0.92	0.92	0.92	0.92	0.92
Canals and Ditches	0.03	0.03	0.03	0.03	0.03	0.03	0.03
CO₂	0.01	+	+	+	+	+	+
Reservoirs	0.01	+	+	+	+	+	+
Constructed Ponds	+	+	+	+	+	+	+
Net Emissions	2.10	2.09	2.09	2.09	2.09	2.09	2.09

“+” positive values less than 0.01mmt

Cropland and Grassland

Cropland, grassland, and settlement areas are defined using the USDA’s National Resources Inventory (NRI) (USDA 2020b) and the National Land Cover Dataset (USGS 2000). Land converted to cropland includes all cropland that was in another land use category within the previous 20 years. The same timeframe applies to land converted to grassland. Estimates of annual CO₂ sequestration in soil organic carbon on cropland remaining cropland and land converted to cropland were made based on soil organic carbon stock changes modeled with the Daycent biogeochemical model (EPA 2023a). Stock changes in biomass, litter and dead organic matter on cropland are not counted but are included for other land use categories (e.g., forest land) that are converted to cropland. Emissions resulting from land converted to cropland more

than offset the annual CO₂ sequestration by the soil of cropland remaining cropland (Table SR3.16). Conversions of forest land to cropland dominated these emissions, averaging 1.3mmt CO₂ per year from 2005 to 2021 (Walters et al. 2023).

Table SR3.16 Cropland, Grassland, and Settlement Net Emissions and Removals, 1990-2021 (mmt CO₂)

Land Use Change	1990	2005	2017	2018	2019	2020	2021
Cropland Remaining Cropland	-1.20	-0.76	-0.83	-0.75	-0.71	-0.80	-0.75
Land Converted to Cropland	1.10	0.83	1.29	1.29	1.29	1.28	1.29
Grassland Remaining Grassland	-0.14	-0.32	-0.05	-0.05	-0.03	-0.09	-0.06
Land Converted to Grassland	-0.24	-0.66	-0.22	-0.21	-0.21	-0.22	-0.22
Settlement Remaining Settlement	0.42	0.42	0.47	0.47	0.47	0.47	0.47
Land Converted to Settlement	1.25	1.27	1.27	1.27	1.27	1.27	1.27
Settlement Trees	-3.52	-3.91	-4.09	-4.08	-4.08	-4.26	-4.28
Net Emission Removals	-2.32	-3.13	-2.15	-2.06	-1.99	-2.34	-2.28

"-" negative values greater than 0.01mmt

Settlements

Settlement trees are located in settlements that remain settlements and other land use categories that are converted to settlements. They contributed a significant amount to annual carbon sequestration (-4.28mmt CO₂ in 2021). However, conversion of forest land to settlements is also a large source of CO₂ as stored forest carbon is released to the atmosphere (1.36mmt CO₂ in 2021 and an average of 1.36mmt CO₂ per year since 2005 (Walters et al. 2023)). Emissions associated with settlements remaining settlements are produced by organic soils in these areas, which are assumed to emit CO₂ at a similar rate to cropland on drained organic soil (EPA 2023a).

Planned Improvements

Improvements to the AFOLU sector inventory are ongoing. Like the “key categories” prioritized by the IPCC, certain areas of improvement will be prioritized here because they are expected to have a significant influence on statewide emission calculations.

Forest Lands, Settlement Trees, and Harvested Wood Products

Future inventory reports may incorporate forest land-cover maps based on satellite images that are available from the National Land Cover Dataset (NLCD). The NLCD land-cover maps incorporate CCAP land-cover data and should therefore reduce any overlap or gaps that currently exist between the inventoried forest land and mapped wetlands land cover. LiDAR data may be used when and where they are available to estimate aboveground biomass in forest land. Comparisons between LiDAR-based biomass estimates and the Forest Service’s field plot measurements may be used to extrapolate coverage.

Expanded collection of survey data from sawtimber producers and manufacturing data for other wood products (for example, paper and particle board), as well as analyses of recent historical trends may be incorporated into future inventories. Product-specific models of emissions as HWPs decay in landfills may also be developed to account for the impact that these products have once they have exceeded their useful lives.

Soil Organic Carbon emissions

Future inventories may incorporate calculations for soil organic carbon emissions from agricultural mineral soils and drained Histosols based on the IPCC (2019) to have a more consistent and localized estimation of SOC emissions than is currently utilized from national data.

Croplands and Grasslands

Future work will specify how soil carbon stock changes are affected by tillage, crop types, and crop residue management practices, either in Daycent model results or by using literature-based or IPCC-recommended factors (IPCC 2019).

Wetlands and Other Water Bodies

Improvements to the wetland CH₄ emission factors and CO₂ removal estimates are planned for future inventories. Field research undertaken this year and over the next three years by Meredith Holgerson (Cornell University, College of Agriculture and Life Sciences) will directly measure CH₄ emission factors that are specific to the state’s freshwater wetland ecosystem types. These will be paired with field measurements of CO₂ sequestration in wetland soils and sediment organic carbon to replace estimates made using IPCC default emission factors. Other planned improvements include revising the spatial boundaries that separate wetlands that emit CH₄ from those that do not using regional salt water interface maps. Although the IPCC default threshold is 18 ppt (Section 4.3.1.2 in IPCC 2014), a salinity threshold of 0.5 ppt is incorporated into the CCAP land-cover maps used here. The lower salinity threshold means that this inventory does not capture any CH₄ emissions from wetlands inundated by water with a salinity higher than 0.5 ppt. Holmquist et al. (2018) provide probability distributions of wetland CH₄ emission across salinity gradients that can then be paired with salinity maps to improve

emissions estimates. Finally, constructed ponds, such as farm ponds, are like wetlands in that they potentially sequester CO₂ as organic carbon in sediment and also emit CH₄. Additional work undertaken through the Cornell University project will measure CO₂ and CH₄ emissions and CO₂ sequestration at a range of different constructed ponds. The goal is to identify the net contribution of their GHG emissions and removals and the impacts of different land use and soil management regimes on this contribution.

Abbreviations

AFOLU	Agriculture, Forestry, and Other Land Use
CCAP	NOAA Coastal Change Analysis Program
CH ₄	Methane
CLCPA	NYS Climate Leadership and Community Protection Act
CO ₂	Carbon dioxide
CO ₂ e	Carbon dioxide equivalent
DEC	NYS Department of Environmental Conservation
DHS	U.S. Department of Homeland Security
DMI	Dry matter intake
ECL	Environmental Conservation Law
EPA	U.S. Environmental Protection Agency
FIA	Forest Inventory and Analysis
GHG	Greenhouse gas
GWP	Global Warming Potential
GWP100	100-Year Global Warming Potential
GWP20	20-Year Global Warming Potential
HWP	Harvested wood product
IPCC	Intergovernmental Panel on Climate Change
LiDAR	Light Detection and Ranging
MMS	Manure Management System
mmt	Million metric tons
mt	Metric tons
N	Nitrogen
N ₂ O	Nitrous oxide
NA	Not applicable
NAIP	USDA National Agriculture Imagery Program
NASS	USDA National Agriculture Statistics Service
NE	Not estimated
NFI	National forest inventory
NH ₃	Ammonia
NHD	USGS National Hydrography Dataset
NLCD	National Land Cover Dataset
NOAA	National Oceanic and Atmospheric Administration
NOx	Nitrogen oxides
NRI	National Resources Inventory
NYCRS	New York State Crop Reporting Service
NYCRR	New York Codes, Rules and Regulations

ppt	Parts per thousand
SIT	EPA State Inventory Tool
USDA	United States Department of Agriculture
USFS	United States Forest Service
USGS	United States Geological Survey

References

- AAPFCO. 2008 through 2017. Commercial Fertilizers: 2008-2015. Association of American Plant Food Control Officials. Columbia, MO, USA: University of Missouri.
- Armentano, T. V. 1979. The role of organic soils in the world carbon cycle--problem analysis and research needs. The Institute of Ecology, Indianapolis, Indiana, Produced for the US Department of Energy.
- Baker, M. 2016. Status and Trends: Biosolids Management in New York State. ClearWaters. New York Water Environment Association, Inc.
- Brakebill, J.W. and J.M. Gronberg. 2017. County-level estimates of nitrogen and phosphorus from commercial fertilizer for the conterminous United States, 1987-2012: U.S. Geological Survey data release. <https://doi.org/10.5066/F7H41PKX>.
- Bureau of the Census. 1989. 1987 Census of Agriculture. Volume 1, Geographic Area Series Part 32, New York State and County Data. U.S. Department Commerce, Bureau of the Census.
- Coulston, J.W., D. N. Wear, and J.M. Vose. 2015. "Complex forest dynamics indicate potential for slowing carbon accumulation in the southeastern United States." *Scientific Reports*. 5, 8002. <https://doi.org/10.1038/srep08002>.
- Crooks, S., A.E. Sutton-Grier, T.G. Troxler, N. Herold, B. Bernal, L. Schile-Beers, T. Wirth. 2018. "Coastal wetland management as a contribution to the U.S. National Greenhouse Gas Inventory." *Nature Climate Change*. 8, 1109-1112. <https://doi.org/10.1038/s41558-018-0345-0>.
- DHS. 2022. Navigable waterway network lines in the Homeland Infrastructure Foundation-Level Data <https://hifld-geoplatform.opendata.arcgis.com/>.
- Domke, G.M., C.H. Perry, B.F. Walters, L.E. Nave, C.W. Woodall, C.W. Swanston. "Toward inventory-based estimates of soil organic carbon in forest of the United States." *Ecological Applications*. 27(4), 1223-1235. <https://doi.org/10.1002/eap.1516>.
- Duxbury, J. M. and J. H. Peverly (1978). Nitrogen And Phosphorus Losses From Organic Soils. *Journal Of Environmental Quality* 7(4): 566-570. Domke, G.M.; B.F. Walters, C.L. Giebink, E.J. Greenfield, J.E. Smith, M.C. Nichols, J.A. Knott, S.M. Ogle, J.W. Coulston, J. Steller. 2023. Greenhouse gas emissions and removals from forest land, woodlands, urban trees, and harvested wood products in the United States, 1990-2021. Resour. Bull. WO-101. Washington, DC: U.S. Department of Agriculture, Forest Service, Washington Office. 10 p. <https://doi.org/10.2737/WO-RB-101>.
- Elder, J. W. and R. Lal. 2008. Tillage effects on gaseous emissions from an intensively farmed organic soil in North Central Ohio. *Soil and Tillage Research* 98(1): 45-55.

EPA. 2023. State Greenhouse Gas Inventory and Projection Tool. June 2023. Washington, D.C.: U.S. Environmental Protection Agency. <https://www.epa.gov/statelocalenergy/download-state-inventory-and-projection-tool>

EPA. 2023a. Inventory of U.S. Greenhouse Gas Emissions and Sinks, 1990-2021. EPA 430-R-23-002. Washington, D.C.: U.S. Environmental Protection Agency.

EPA. 2023b. Methodology Report for Inventory of U.S. Greenhouse Gas Emissions and Sinks by State: 1990-2021. EPA-430-R-23-003. Washington, D.C.: U.S. Environmental Protection Agency.

EPA. 2023c. National Lakes Assessment 2017: Technical Support Document. EPA-841-R-22-001. Washington, DC: Office of Water and Office of Research and Development. <https://www.epa.gov/national-aquatic-resource-surveys/national-lakes-assessment-2017-technical-support-document>

Holgerson, M.A. and P.A. Raymond. 2016. "Large contribution to inland water CO₂ and CH₄ emissions from very small ponds." *Nature Geoscience*. 9, 222-226. <https://doi.org/10.1038/NGEO2654>.

Holmquist, J.R., L. Windham-Myers, N. Bliss, S. Crooks, J.T. Morris, P. Megonigal, T. Troxler, D. Weller, et al. 2018. "Accuracy and precision of tidal wetland soil carbon mapping in the conterminous United States." *Scientific Reports*. 8, 9478. <https://doi.org/10.1038/s41598-018-26948-7>.

IPCC. 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories. [H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe (eds.)] Hayama, Japan: The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change.

IPCC. 2013. Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. [Stocker, T.F., D. Qin, G.-K., Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press.

IPCC. 2014. 2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands. [Hiraishi, T., T. Krug, K. Tanabe, N. Srivastava, J. Baasansuren, M. Fukuda, T.G. Troxler (eds.)]. The National Greenhouse Gas Inventories Technical Support Unit, The Intergovernmental Panel on Climate Change.

IPCC. 2019. 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories. [Calvo Buendia, E., Tanabe, K., Kranjc, A., Baasansuren, J., Fukuda, M., Ngarize, S., Osako, A., Pyrozhenko, Y., Shermanau, P. and Federici, S. (eds.)]. Hayama, Kanagawa, Japan: The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change.

Neubauer, S. 2014. "On the challenges of modeling the net radiative forcing of wetlands: reconsidering Mitsch et al. 2013." *Landscape Ecology*. 29, 571-577. <https://doi.org/10.1007/s10980-014-9986-1>.

NYSDEC. 1999. Solid Waste Facts. New York State Department of Environmental Conservation, Division of Materials Management. 1-20.

NYSDEC. 2018. Biosolids Management in New York State. New York State Department of Environmental Conservation, Division of Materials Management, Albany, NY.

NYS Seagrass Taskforce. 2009. Final report of the New York State Seagrass Task Force: Recommendations to the New York State governor and legislature. New York State Department of Environmental Conservation. https://www.dec.ny.gov/docs/fish_marine_pdf/finalseagrassreport.pdf

Oswalt, S.N., W.B. Smith, P.D. Miles, S.A. Pugh. 2019. Forest Resources of the United States, 2017: a technical document supporting the Forest Service 2020 RPA Assessment. Gen. Tech. Rep. WO-97. U.S. Department of Agriculture, Forest Service. <https://doi.org/10.2737/WO-GTR-97>.

Ropel, S.C. and B.L. Smith. 2001. New York Agricultural Statistics 2000-2001. New York Agricultural Statistics Service. https://www.nass.usda.gov/Statistics_by_State/New_York/Publications/Annual_Statistical_Bulletin/2001/01-complete.pdf

Salinas, C., C.M. Duarte, P.S. Lavery, P. Masque, A. Arias-Ortiz, J.X. Leon, D. Callaghan, G.A. Kendrick, O. Serrano. 2020. "Seagrass losses since mid-20th century fuelled CO₂ emissions from soil carbon stocks." *Global Change Biology*. 26, 4772-4784. <https://doi.org/10.1111/gcb.15204>.

USACE. 2021. National Inventory of Dams. <https://nid.sec.usace.army.mil/>

USDA. 2008. Muckland Report September 2008. New York Crop Reporting Service. Albany, NY: New York Agricultural Statistics Service, U.S. Department of Agriculture.

USDA. 2019. 2017 Census of Agriculture United States Summary and State Data. AC-17-A-51. Washington, D.C.: U.S. Department of Agriculture.

USDA. 2020a. Quickstats. U.S. Department of Agriculture National Agricultural Statistics Service. https://www.nass.usda.gov/Quick_Stats/.

USDA. 2020b. Summary Report: 2017 National Resources Inventory. Washington, DC and Ames Iowa: Natural Resources Conservation Service and Center for Survey Statistics and Methodology, Iowa State University.

USGS. 2000. National Land Cover Dataset. <https://doi.org/10.3133/fs10800>

USGS. 2016. National Hydrography Dataset (NHD) Downloadable Data Collection – National Geospatial Data Asset (NGDA). Rolla, MO and Denver, CO: National Geospatial Technical Operations Center. <http://nhd.usgs.gov>.

USGS. 2018. Crushed Stone, 2016 Minerals Yearbook. U.S. Department of the Interior/U.S. Geological Survey, Washington, D.C.

Walters, B.F., G.M. Domke, E.J. Greenfield, J.E. Smith, S.M. Ogle. 2023. Greenhouse gas emissions and removals from forest land, woodlands, and urban trees in the United States, 1990-2021: Estimates and quantitative uncertainty for individual states, regional ownership groups, and National Forest System regions. Fort Collins, CO: Forest Service Research Data Archive. <https://doi.org/10.2737/RDS-2023-0020>

West, T.O. and A.C. McBride. 2005. “The contribution of agricultural lime to carbon dioxide emissions in the United States: dissolution, transport, and net emissions.” *Agriculture Ecosystems & Environment* 108:145-154. <https://doi.org/10.1016/j.agee.2005.01.002>.

Westfall, J.A., Coulston, J.W., Gray, A.N., Shaw, J.D., Radtke, P.J., Walker, D.M., Weiskittel, A.R., MacFarlane, D.W., Affleck, D.L.R., Zhao, D.T., Hailemariam, P., Krishna, P., Frank, J.M., Prisley, S.P., Wang, Y., Sánchez Meador, A.J., Auty, D., Domke, G.M. [In press]. “A national-scale tree volume, biomass, and carbon modeling system for the United States, . Gen., . Tech. Rep. WO-104. Washington, DC: U.S. Department of Agriculture, Forest Service. 60 p. <https://doi.org/10.2737/WO-GTR-104>.